arXiv:1205.3274v2 [math.NT] 12 Jul 2012

LOWER BOUNDS ON THE ARITHMETIC SELF-INTERSECTION NUMBER OF THE RELATIVE DUALIZING SHEAF ON ARITHMETIC SURFACES

ULF KÜHN, JAN STEFFEN MÜLLER

Abstract. We provide an explicitly computable lower bound for the arithmetic self-intersection number $\overline{\omega}^2$ of the dualizing sheaf on a large class of arithmetic surfaces. If some technical conditions are satisfied, then this lower bound is positive. In particular, these technical conditions are always satisfied for minimal arithmetic surfaces with at least one reducible fiber and without multiple components.

1. Introduction

Let k be a number field, let $\mathcal{O}_k$ denote the ring of integers of k. Let X/k denote a smooth, projective, geometrically irreducible curve of genus $g > 1$. Finally, let $\pi : \mathcal{X} \to \operatorname{Spec}(\mathcal{O}_k)$ be a proper regular model of X.

Let $\overline{\omega} = \overline{\omega}_{\mathcal{X}}$ denote the relative dualizing sheaf on $\mathcal{X}$ over $\operatorname{Spec}(\mathcal{O}_k)$, equipped with the Arakelov metric. The arithmetic self-intersection $\overline{\omega}^2$ is one of the most fundamental objects in arithmetic intersection theory. In this note we show how it is possible to compute lower bounds on $\overline{\omega}^2$ under a technical hypothesis.

Our main result Theorem 1.4 implies the following:

Theorem 1.1. *If $\mathcal{X}$ is minimal and all irreducible components of $\mathcal{X}$ have multiplicity 1, then there is an effectively computable lower bound* (1) *on $\overline{\omega}^2$. Moreover, this lower bound is always nonnegative and is zero if and only if all fibers of $\mathcal{X}$ are irreducible.*

As an immediate corollary we recover the following previously known result from [17], [13] and [4]. Here we call a fiber *reducible* if it has more than one irreducible component.

Corollary 1.2. *If $\mathcal{X}$ is semistable and minimal and has at least one reducible fiber, then there is an effectively computable positive lower bound on $\overline{\omega}^2$.*

The positivity of $\overline{\omega}^2$ has been shown without assuming semistability, but still assuming the existence of at least one reducible fiber in [14]. However, this approach is not suitable for explicit computations of such bounds.

To each $\mathbb{Q}$-divisor $D \in \operatorname{Div}_{\mathbb{Q}}(X) := \operatorname{Div}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$ of degree one we attach in Definiton 3.1 a hermitian line bundle $\overline{L_D}$. For the construction we use vertical divisors V_D and U_D (see (6)) satisfying certain properties whose existence is guaranteed by Proposition 2.1. This line bundle is our main tool in what follows. Namely, we prove that if we take $D = K$, where $(2g-2)K$ is a canonical divisor, then showing $\overline{L_K}^2 \geq 0$ implies our lower bound.

By construction, the height function $h_{\overline{L_D}}(\,\cdot\,)$ defined in (8) on the curve X is closely related to the Néron-Tate height induced by an embedding of X into its Jacobian via D. In particular,

---

Date: March 4, 2013.

The second author was supported by DFG-grant KU 2359/2-1.

we can show $h_{\overline{L_D}}(P) = h_{\mathrm{NT}}([P - D])$ for any $P \in X(k)$, whenever the Zariski closure $D_{\mathcal{X}}$ of D in $\mathcal{X}$ is disjoint from a certain finite set $T(\mathcal{X}) \subset \mathcal{X}$ of closed points (cf. Definition 4.2). If $E \in \mathrm{Div}_{\mathbb{Q}}(X)$ is effective such that the support of its Zariski closure $E_{\mathcal{X}} \in \mathrm{Div}_{\mathbb{Q}}(\mathcal{X})$ is also disjoint from $T(\mathcal{X})$, then $h_{\overline{L_D}}(E) \geq 0$ (cf. Proposition 4.6). For such D a refinement of Zhang's results on positive hermitian line bundles [16, Theorem 6.5] allows us to prove the inequality $\overline{L_D}^2 \geq 0$; more precisely we show in Section 5:

Theorem 1.3. *If $\overline{L_D}$ is relatively semipositive and $D_{\mathcal{X}} \cap T(\mathcal{X}) = \emptyset$, then we have $\overline{L_D}^2 \geq 0$.*

Here a line bundle is *relatively semipositive* if its restriction to any irreducible vertical divisor has nonnegative arithmetic degree (see Definition 3.2).

Theorem 1.3 is used in Section 6 to prove our main result:

Theorem 1.4. *Let $K \in \mathrm{Div}_{\mathbb{Q}}(X)$ be a $\mathbb{Q}$-divisor such that $(2g-2)\cdot K$ is a canonical $\mathbb{Q}$-divisor on X and such that the Zariski closure $K_{\mathcal{X}}$ of K on $\mathcal{X}$ satisfies $K_{\mathcal{X}} \cap T(\mathcal{X}) = \emptyset$.*

If the hermitian line bundle $\overline{L_K}$ is relatively semipositive then we have

$$\overline{\omega}^2 \geq -\frac{g-1}{g}\mathcal{O}(2V_K + U_K)^2 + 2(\overline{\omega}\,.\,\mathcal{O}(U_K)). \tag{1}$$

We would like to stress that it is easy to compute the divisors V_K and U_K; hence the right hand side of (1) can be computed easily for a given arithmetic surface.

If all irreducible components of $\mathcal{X}$ have multiplicity one, then we show in Proposition 3.4 that $\overline{L_D}$ is relatively semipositive for all D of degree one. In that case (1) gives in fact a lower bound, since there always exists a K satisfying $K_{\mathcal{X}} \cap T(\mathcal{X}) = \emptyset$. A priori that lower bound could be negative, however with the additional asumption of minimality of $\mathcal{X}$, we derive Theorem 1.1 from Theorem 1.4 in Section 5.

We believe that for all minimal arithmetic surfaces there exists a relatively semipositive $\overline{L_K}$ such that (1) is a nonnegative lower bound on $\overline{\omega}^2$. Indeed, if there are components of higher multiplicities, one can still check whether Theorem 1.4 is applicable. As an example, we prove that the conditions of Theorem 1.4 are satisfied for the minimal regular model $\mathcal{F}_p^{\min}$ of the Fermat curve of prime exponent $p > 3$ over the p-th cyclotomic numbers and that the resulting lower bound is positive. This does not follow from Theorem 1.1, since the irreducible components of $\mathcal{F}_p^{\min}$ need not have multiplicity one.

Theorem 1.5. *The arithmetic self-intersection $\overline{\omega}^2$ of the relative dualizing sheaf on $\mathcal{F}_p^{\min}$ satisfies*

$$\overline{\omega}^2 \geq \frac{1}{4p^3(p-1)(p-2)}\left(4p^6 - 32p^5 + \frac{13}{2}p^4 + \frac{73}{2}p^3 - 52p^2 + 144\right)\log p.$$

This lower bound is positive for all $p > 7$. Furthermore, if $p = 5$, then we have $\overline{\omega}^2 \geq \frac{188}{125}\log 5$ and if $p = 7$, then we have $\overline{\omega}^2 \geq \frac{37277}{6860}\log 7$.

In Section 7 we use Theorem 1.4 to prove Theorem 1.5. We then compare the resulting lower bound to the upper bound computed by Curilla and Kühn in [7].

We have no explicit result how sharp the bound (1) is. An upper bound on $\overline{L_K}^2$ can be obtained from [16, Theorem 6.5], which implies that

$$\overline{L_K}^2 \leq 4ge'_{\overline{L_K}},$$

where $e'_{\overline{L_K}} = \liminf_{P \in X(\bar{k})} h_{\overline{L_K}}(P)$. Hence we find

$$0 \leq \overline{\omega}^2 - \left(-\frac{g-1}{g} \mathcal{O}(2V_K + U_K)^2 + 2(\overline{\omega} \,.\, \mathcal{O}(U_K)) \right) \leq 4(g-1) e'_{\overline{L_K}}, \tag{2}$$

so $e'_{\overline{L_K}}$ controls how sharp our bound is.

For the rest of the introduction we assume that $\mathcal{X}$ is semistable. In that case lower bounds on $\overline{\omega}^2$ can also be derived by means of the admissible intersection theory due to Zhang, cf. [17, Theorem 5.5]. Namely, Zhang shows that

$$\overline{\omega}^2 = \omega_a^2 + r,$$

where ω_a^2 is the admissible self-intersection of ω and r is some explicitely computable invariant. It is given in terms of admissible Green's functions on the reduction graphs of the special fibers of $\mathcal{X}$. Furthermore, he deduces Corollary 1.2 from the inequality $\omega_a^2 \geq 0$ and the fact that $r > 0$ if there exists a fiber which does not have potential good reduction. In fact he proves, similarly to (2),

$$0 \leq \overline{\omega}^2 - r \leq 4(g-1) a'(K),$$

where $a'(K) = \liminf_{P \in X(\bar{k})} h_{\mathrm{NT}}([P-K])$ (see[17, Theorem 5.6]).

Zhang's approach has been employed by Abbes-Ullmo [1] and Mayer [11] to determine lower bounds for certain modular curves. Note that this approach does not apply to non-semistable arithmetic surfaces, so in particular it cannot be used for $\mathcal{F}_p^{\min}$.

The positivity of $\overline{\omega}_a^2$ is equivalent to the Bogomolov conjecture for the Jacobian of X, which was proved by Ullmo [15]. An explicit strictly positive lower bound for $\overline{\omega}_a^2$ would imply an effective version of the Bogomolov conjecture for X by [16, Theorem 5.6]. We point out that for $\mathcal{X}$ with at least one bad fiber our lower bound (1) minus Zhang's quantity r is a lower bound for $\overline{\omega}_a^2$.

We would like to thank Ariyan Javanpeykar and David Holmes for a careful reading of the manuscript and many helpful suggestions. We also thank Zubeyir Cinkir for advice on the proof of Proposition 3.4 and Christian Curilla for drawing Figure 1.

2. Correction divisors

First we introduce some notation which will be used in several places throughout this work. If v is a non-archimedean place of k and $\mathcal{X}_v = \sum_{i=1}^{r_v} b_i \Gamma_i$, where $\{\Gamma_1, \ldots, \Gamma_{r_v}\}$ is the set of irreducible components of $\mathcal{X}_v$, then we define an $r_v \times r_v$ matrix $M_{\mathcal{X}_v} = (m_{ij})_{i,j}$ as minus the intersection matrix of $\mathcal{X}_v$, that is $m_{ij} = -(b_i \Gamma_i \,.\, b_j \Gamma_j)$. Let $M^+_{\mathcal{X}_v} = (n_{ij})_{i,j}$ denote the Moore-Penrose pseudoinverse of M. Then both $M_{\mathcal{X}_v}$ and $M^+_{\mathcal{X}_v}$ are symmetric and positive semidefinite. Moreover, the following facts hold (see [6]):

$$\sum_j m_{ij} = \sum_j n_{ij} = 0 \quad \text{for all } i \in \{1, \ldots, r_v\}, \tag{3}$$

$$\sum_{j=1}^{r} n_{kj} m_{ji} = -\frac{1}{r_v} + \delta_{ik} \text{ for all } i, k \in \{1, \ldots, r_v\} \tag{4}$$

We now associate a metrized graph (see [5] or [17]) $\mathcal{G}_{\mathcal{X}_v}$ to $\mathcal{X}_v$ as follows: The vertex set $V(\mathcal{G}_{\mathcal{X}_v})$ is given by $\{\Gamma_1, \ldots, \Gamma_{r_v}\}$ and two vertices Γ_i and Γ_j are connected by a single edge of length $-1/m_{ij}$ if $m_{ij} \neq 0$. Then $M_{\mathcal{X}_v}$ is the discrete Laplacian associated to $\mathcal{G}_{\mathcal{X}_v}$ (see [6]).

For the remainder of this section we work over the strict henselization k_v^{nr} of the completion of k at v and its ring of integers $\mathcal{O}_v^{\mathrm{nr}}$. If $E \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$, then we let $v_v(E)$ be the column vector with entries $v_{v,i}(E)$, where

$$v_{v,i}(E) = (b_i\Gamma_i\,.\,E_{\mathcal{X}}).$$

Moreover, we set for each $i \in \{1, \ldots, r_v\}$

$$a_{v,i} := -\Gamma_i^2 + 2p_a(\Gamma_i) - 2,$$

where $p_a(\Gamma_i)$ is the arithmetic genus of Γ_i,

$$a'_{v,i} := \frac{1}{2g-2} a_{v,i}$$

and

$$w_{v,i}(E) = \frac{1}{\deg(E)}(b_i \cdot a_{v,i} - v_{v,i}(E)).$$

Proposition 2.1. *For any $E \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ of degree d and any non-archimedean place v of k there exists a vertical $\mathbb{Q}$-divisor $V_v(E)$ such that*

$$(E_{\mathcal{X}} + V_v(E)\,.\,\Gamma) = \frac{d}{2g-2}\left(-\Gamma^2 + 2p_a(\Gamma) - 2\right)$$

for every irreducible vertical component Γ of $\mathcal{X}_v$, where $p_a(\Gamma)$ is the arithmetic genus of Γ.

Proof. We first assume that $d = 2g-2$ and that E is effective. Let v be a non-archimedean place of k and let $M = M_{\mathcal{X}_v}$. Then a divisor

$$V_v(E) = \sum_{i=1}^{r_v} b_i c_i(E)\Gamma_i$$

as desired exists if and only if the system

$$-M \cdot c(E) = w(E) \tag{5}$$

of linear equations has a solution, where $c(E)$ is the column vector with entries $c_i(E)$ and $w(E)$ is the column vector with entries $w_i(E)$. Note that (5) is solvable if and only if the rank of $(-M|w(E))$ is equal to $\mathrm{rk}(-M) = r_v - 1$, which is equivalent to $\sum_{i=1}^{r_v} w_i(E) = 0$ because of (3). But it is easy to see that $\sum_{i=1}^{r_v} w_i(E)$ only depends on the degree of E. It is shown in [7] that (5) is solvable if E is a canonical $\mathbb{Q}$-divisor on $X \times k_v^{\mathrm{nr}}$, so we get the existence of a solution for all E of degree $2g-2$.

Next let E be effective of arbitrary degree $d > 0$. Then we set

$$V_v(E) = \frac{d}{2g-2} V_v\left(\frac{2g-2}{d}E\right).$$

Finally, if E is not effective, then we write $E = E_1 - E_2$, where E_1 and E_2 are effective and set $V_v(E) = V_v(E_1) - V_v(E_2)$. It is obvious that $V_v(E)$ does not depend on the choice of E_1 and E_2. □

Proposition 3 says that we can extend E to a divisor on $\mathcal{X} \times \mathcal{O}_v^{\mathrm{nr}}$ which satisfies the adjunction formula up to a factor $\deg(E)$ on the special fiber.

The case of degree zero is especially interesting. It is well-known (see for instance [9, Theorem III.3.6]) that for any $Z \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ of degree zero we can find a vertical $\mathbb{Q}$-divisor $\Phi_{\mathcal{X}}(Z) = \sum_v \Phi_{\mathcal{X},v}(Z)$ on $\mathcal{X} \times \mathcal{O}_v^{\mathrm{nr}}$ such that $Z_{\mathcal{X}} + \Phi_{\mathcal{X}}(Z)$ has trivial intersection with all vertical divisors on $\mathcal{X}_v$.

Corollary 2.2. *Let* $Z = E - E' \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ *have degree zero. Then we can take* $\Phi_{\mathcal{X},v}(Z) = V_v(E) - V_v(E')$. □

Remark 2.3. On a side note, we can define, for each non-archimedean place v of k, a local pairing on coprime divisors $E_1, E_2 \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ by

$$[E_1, E_2]_v := (E_{1,\mathcal{X}} + V_v(E_1) \,.\, E_{2,\mathcal{X}} + V_v(E_2)).$$

Corollary 2.2 implies that $[E_1, E_2]_v$ extends the local Néron pairing (see [9, §III.5]) to divisors of arbitrary degree. To our knowledge, this has not been done before without using the reduction graph as in [5] or [17].

Moreover, if v is archimedean, then we can define $[E_1, E_2]_v$ to be the admissible pairing of E_1 and E_2 at v as in [17, §4.5]. Then we get a global pairing

$$[E_1, E_2] := \sum_v [E_1, E_2]_v$$

on divisors $E_1, E_2 \in \mathrm{Div}_{\mathbb{Q}}(X)$ with disjoint support. This pairing has the following properties, which may be of independent interest:

- (i) $[\cdot, \cdot]$ is bilinear and symmetric.
- (ii) $[E_1, \mathrm{div}(f)] = 0$ for any $f \in k(X)^*$. Hence $[\cdot, \cdot]$ induces a well-defined pairing on divisor classes.
- (iii) If $\deg(E_1) = \deg(E_2) = 0$, then we have $[E_1, E_2] = -(E_1, E_2)_{\mathrm{NT}}$, where the latter is the Néron-Tate height pairing.
- (iv) If E_1 and E_2 are canonical divisors on X, then we have $[E_1, E_2] = \overline{\omega}^2$.

3. THE HERMITIAN LINE BUNDLE $\overline{L_D}$

Throughout this section we let D denote a $\mathbb{Q}$-divisor on X of degree one. We define a vertical $\mathbb{Q}$-divisor U_D on $\mathcal{X}$ as follows: If v is a non-archimedean place of k such that $\Gamma_1^v, \ldots, \Gamma_{r_v}^v$ are the irreducible components of the special fiber $\mathcal{X}_v$ and $i \in \{1, \ldots, r_v\}$, we let $G_i^v \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ be a $\mathbb{Q}$-divisor on X of degree one such that the Zariski closure $G^v_{i,\mathcal{X}\times\mathcal{O}_v^{\mathrm{nr}}}$ intersects Γ_i^v, but does not intersect Γ_j^v for $j \neq i$. It follows from [2, Corollary 9.2.10] that such a $\mathbb{Q}$-divisor always exists. Let $\gamma_{D,v,i} := V_v(G_i^v)^2 - 2(V_v(G_i^v) \,.\, V_v(D \times k_v^{\mathrm{nr}}))$. Then we define $U_{D,v}$ by

$$U_{D,v} = -\sum_{i=1}^{r} \gamma_{D,v,i} \cdot \Gamma_i^v$$

and set

$$U_D = \sum_v U_{D,v}. \tag{6}$$

Definition 3.1. We define a hermitian line bundle $\overline{L_D}$ by

$$\overline{L_D} = \overline{\omega} \otimes \overline{\mathcal{O}}(2D_{\mathcal{X}}) \otimes \mathcal{O}(U_D)^{-1} \otimes \overline{\mathcal{O}}(-a\mathcal{X}_\infty),$$

where $a = \overline{\mathcal{O}}(D_{\mathcal{X}})^2 - \mathcal{O}(V_D)^2 \in \mathbb{R}$. Here $\overline{\mathcal{O}}(D_{\mathcal{X}})^2$ is the self-intersection of $\mathcal{O}(D_{\mathcal{X}})$, equipped with the Arakelov metric, $V_D = \sum_v V_v(D)$ and $\overline{\mathcal{O}}(-a\mathcal{X}_\infty) = (\mathcal{O}_{\mathcal{X}}, |\cdot|e^a)$.

Unless otherwise stated, all metrics will be Arakelov metrics (so that the Arakelov adjunction formula holds, cf. [9, §IV.5]), except for vertical line bundles, which are equipped with the trivial metric throughout.

The hermitian line bundle $\overline{L_D}$ will be our main object of interest. We want to show that it has nonnegative self-intersection.

Definition 3.2. We say that a hermitian line bundle $\overline{L}$ is *relatively semipositive* if it has nonnegative intersection with all irreducible vertical components of $\mathcal{X}$. If $\overline{L}$ has nonnegative (resp. positive) intersection with all irreducible horizontal divisors on $\mathcal{X}$, then we call $\overline{L}$ *horizontally semipositive* (resp. *horizontally positive*).

Next we will show that $\overline{L_D}$ is relatively semipositive, at least when all components of the special fibers of $\mathcal{X}$ have multiplicity one, which we assume for the remainder of this section.

We fix a non-archimedean place v of k, keep the notation of the previous section 2 and let $M_{\mathcal{X}_v} = (m_{ij})_{i,j}$ and $M^{+}_{\mathcal{X}_v} = (n_{ij})_{i,j}$. For each divisor $E \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ we write $V_v(E) = \sum_{i=1}^{r_v} c_{v,i}(E)\Gamma_i$, since by assmuption all $b_i = 1$. For simplicity, we will drop the subscripts v for the remainder of this section.

Lemma 3.3. *We have*

$$-(U_D \,.\, \Gamma_i) = \sum_{j=1}^{r} n_{jj} m_{ij} - 2v_i(D) + \frac{2}{r}$$

Proof. This is an easy computation using the properties of $M_{\mathcal{X}_v}$ listed in Section 2:

$$\begin{aligned}
-(U_D \,.\, \Gamma_i) &= \sum_{l=1}^{r}\sum_{j=1}^{r} c_j(G_l)\cdot(w_j(G_l) - 2w_j(D))\cdot m_{lj}\\
&= \sum_{l=1}^{r}\left(\sum_{j=1}^{r} c_j(G_l)\cdot(2v_j(D) - a'_j - \delta_{lj})\right)\cdot m_{li}\\
&= \sum_{l=1}^{r}\sum_{j=1}^{r}\sum_{k=1}^{r}\big(n_{kj}(a'_k - \delta_{kl})\cdot(2v_j(D) - a'_j - \delta_{lj})\big)\cdot m_{li}\\
&= \sum_{l=1}^{r}\sum_{j=1}^{r}\sum_{k=1}^{r}\big(2n_{kj}a'_k v_j(D) - n_{kj}a'_k a'_j - n_{kj}a'_k\delta_{lj} - 2n_{kj}\delta_{kl}v_j(D)\\
&\qquad\qquad + n_{kj}\delta_{kl}a'_j + n_{kj}\delta_{kl}\delta_{lj}\big)\cdot m_{li}\\
&= \sum_{j=1}^{r} n_{jj}m_{ij} - 2\sum_{l=1}^{r}\sum_{j=1}^{r} n_{lj}v_j(D)m_{li}\\
&= \sum_{j=1}^{r} n_{jj}m_{ij} - 2\sum_{j=1}^{r} v_j(D)\sum_{l=1}^{r} n_{lj}m_{li}\\
&= \sum_{j=1}^{r} n_{jj}m_{ij} - 2\sum_{j=1}^{r} v_j(D)\left(\delta_{ij} - \frac{1}{r}\right)\\
&= \sum_{j=1}^{r} n_{jj}m_{ij} - 2v_i(D) + \frac{2}{r}.
\end{aligned}$$

□

It turns out that in the case of simple multiplicities, Lemma 3.3 suffices to prove relative semipositivity of $\overline{L_D}$.

Proposition 3.4. *Suppose that all components of the fibers of $\mathcal{X}$ have simple multiplicities. Then for any $D \in \mathrm{Div}_{\mathbb{Q}}(X)$ of degree 1 the hermitian line bundle $\overline{L_D}$ is relatively semipositive.*

Proof. Let $i \in \{1, \ldots, r\}$.

$$\begin{aligned}\frac{(\overline{L_D} \,.\, \mathcal{O}(\Gamma_i))}{\log Nm(\mathfrak{p}_v)} &= (\mathcal{K} \,.\, \Gamma_i) + 2(D_{\mathcal{X}} \,.\, \Gamma_i) - (U_{D,v} \,.\, \Gamma_i) \\ &= (\mathcal{K} \,.\, \Gamma_i) + 2v_i(D) + \sum_{j=1}^{r} n_{jj} m_{ij} + \frac{2}{r} - 2v_i(D) \\ &= m_{ii} + 2p_a(\Gamma_i) - 2 + \sum_{j=1}^{r} n_{jj} m_{ij} + \frac{2}{r}.\end{aligned}$$

Here $\mathcal{K}$ is a canonical divisor on $\mathcal{X}$ and $p_a(\Gamma_i)$ is the arithmetic genus of Γ_i. The second equality follows from $\sum_{j=1}^{r} v_j(D) = 1$ and (3) whereas the third equality follows from the adjunction formula on $\mathcal{X}_v$. Note that $(\overline{L_D} \,.\, \mathcal{O}(\Gamma_i))$ is independent of D.

It follows that

$$\begin{aligned}\sum_j n_{jj} m_{ij} &= \sum_j (n_{jj} + n_{ii} - 2n_{ij}) m_{ij} - \frac{2}{r} + 2 \\ &= \sum_j r(\Gamma_i, \Gamma_j) m_{ij} - \frac{2}{r} + 2,\end{aligned}$$

where $r(\Gamma_i, \Gamma_j)$ is the effective resistance between the nodes Γ_i and Γ_j if we consider the metrized graph $\mathcal{G}_{\mathcal{X}_v}$ as a resistive electric circuit as, for instance, in [5, 17]. The formula $r(\Gamma_i, \Gamma_j) = n_{jj} + n_{ii} - 2n_{ij}$ is taken from [6].

In order to prove the proposition it therefore suffices to show that

$$m_{ii} + \sum_j r(\Gamma_i, \Gamma_j) m_{ij} \geq 0. \tag{7}$$

But we can rewrite the left hand side as

$$\sum_{j \neq i} m_{ij} (r(\Gamma_i, \Gamma_j) - 1).$$

A component Γ_j can only contribute a negative summand to the sum if $m_{ij} \neq 0$. This means that the nodes on $\mathcal{G}_{\mathcal{X}_v}$ corresponding to Γ_i and Γ_j are connected by an edge of length $-\frac{1}{m_{ij}} \leq 1$. In this case it is well known that the effective resistance $r(\Gamma_i, \Gamma_j)$ is bounded from above by the edge length. Hence all terms in the sum are nonnegative, proving (7) and thus the proposition. □

If we want to prove relative semipositivity for arbitrary arithmetic surfaces, estimates are more difficult. Nevertheless, we expect Proposition 3.4 to hold for all minimal arithmetic surfaces. See Lemma 7.3 for a family of examples.

4. Heights and intersections

If ℓ is a finite extension of k, then we let $\varphi_\ell := \mathrm{pr}_1 \circ \pi_\ell : \mathcal{X}^\ell \to \mathcal{X}$, where $\pi_\ell : \mathcal{X}^\ell \to \mathcal{X} \times \mathcal{O}_\ell$ denotes the minimal desingularization. For the definition of semistable arithmetic surfaces we refer to 4.1; in particular, we do not require a semistable arithmetic surface to be minimal.

Lemma 4.1. *There exists a finite extension ℓ_0/k such that $\mathcal{X}^\ell$ is semistable for any finite extension ℓ/ℓ_0.*

Definition 4.2. Let ℓ_0/k be as in Lemma 4.1. We denote the smooth locus of $\mathcal{X}^{\ell_0}$ by $\mathcal{X}^{\ell_0}_{\mathrm{sm}}$ and we denote the exceptional locus of φ_{ℓ_0} by $\mathrm{Exc}(\varphi_{\ell_0})$. With this notation we define

$$T(\mathcal{X}) := \varphi_{\ell_0}\left(\mathcal{X}^{\ell_0} \setminus \mathcal{X}^{\ell_0}_{\mathrm{sm}} \, \cup \, \mathrm{Exc}(\varphi_{\ell_0})\right).$$

Remark 4.3. If $\mathcal{X}$ is semistable, then we have $T(\mathcal{X}) = \mathcal{X}^{\ell_0} \setminus \mathcal{X}_{\mathrm{sm}}$.

Lemma 4.4. *Any irreducible divisor E on X such that* $\mathrm{supp}(E_{\mathcal{X}}) \cap T(\mathcal{X}) = \emptyset$ *satisfies $\varphi_\ell^* E_{\mathcal{X}} = E_{\mathcal{X}^\ell}$ for any finite extension ℓ/ℓ_0.*

Proof. Let E be an irreducible divisor on X whose closure $E_{\mathcal{X}}$ does not contain an element of $T(\mathcal{X})$ in its support and let ℓ/k be a finite extension containing ℓ_0. Note that if $\varphi^* E_{\mathcal{X}} \neq E_{\mathcal{X}^\ell}$, then there is an irreducible component $\Gamma \in \mathrm{Exc}(\varphi_\ell)$ such that $\varphi_\ell(\Gamma) \in \mathrm{supp}(E_{\mathcal{X}})$.

But this means that either $\Gamma \in \mathrm{Exc}(\varphi_{\ell_0})$, implying that $\varphi_\ell(\Gamma) \in T(\mathcal{X})$ by definition, or Γ is contracted to a point by the desingularization morphism $\mathcal{X}_\ell \to \mathcal{X}_{\ell_0} \times \mathcal{O}_\ell$. In this case Γ maps to a singular point of $\mathcal{X}_{\ell_0} \times \mathcal{O}_\ell$, whence $\varphi_\ell(\Gamma) \in T(\mathcal{X})$.

□

Let $\mathcal{E}$ be an irreducible horizontal divisor on $\mathcal{X}$. The height of $\mathcal{E}$ with respect to a hermitian line bundle $\overline{L}$ on $\mathcal{X}$ is defined by

$$h_{\overline{L}}(\mathcal{E}) = \frac{(\overline{L} \,.\, \overline{\mathcal{O}}(\mathcal{E}))}{[k:\mathbb{Q}] \deg(\mathcal{E}_k)}, \tag{8}$$

where the metric on $\mathcal{O}(\mathcal{E})$ is admissible in the Arakelov-theoretic sense. We extend this to arbitrary effective divisors on $\mathcal{X}$ by linearity. If E is an irreducible effective divisor on X, then we set

$$h_{\overline{L}}(E) := h_{\overline{L}}(E_{\mathcal{X}}).$$

Definition 4.5. Let T be a finite set of closed points of X. We say that a hermitian line bundle $\overline{L}$ is T*-semipositive* if we have $h_{\overline{L}}(\mathcal{E}) \geq 0$ for all effective horizontal divisors $\mathcal{E}$ on $\mathcal{X}$ such that $\mathrm{supp}(\mathcal{E}) \cap T = \emptyset$.

Let h_{NT} denote the Néron-Tate height on the Jacobian J of X with respect to the symmetrized theta divisor $\Theta + [-1]^*\Theta$. Finally, for each divisor $D \in \mathrm{Div}_{\mathbb{Q}}(X)$ of degree one, let $j_D : X \hookrightarrow J$ be the embedding mapping a point $Q \in X$ to the class of $Q - D$.

Proposition 4.6. *Suppose that $D \in \mathrm{Div}_{\mathbb{Q}}(X)$ has degree one and* $\mathrm{supp}(D_{\mathcal{X}}) \cap T(\mathcal{X}) = \emptyset$. *Then $\overline{L_D}$ is $T(\mathcal{X})$-semipositive. Moreover, if $P \in X(k)$, then $h_{\overline{L_D}}(P) = h_{\mathrm{NT}}(j_D(P))$.*

Proof. Let E be an irreducible divisor on X such that $\mathrm{supp}(E_{\mathcal{X}}) \cap T(\mathcal{X}) = \emptyset$. Let ℓ_0 be as in Lemma 4.1 and let ℓ/k be a finite extension containing ℓ_0 such that E has pointwise ℓ-rational support. Let $d = \deg(E)$ and $E = \sum_{j=1}^{d}(P_j)$, where $P_j \in X(\ell)$. By Proposition 4.4 we have

$$\varphi_\ell^* E_{\mathcal{X}} = E_{\mathcal{X}^\ell} \text{ and } \varphi_\ell^* D_{\mathcal{X}} = D_{\mathcal{X}^\ell}. \tag{9}$$

Hence we get, using [9, Theorem III.4.5],

$$\Phi_{\mathcal{X}^\ell}(E - dD) = \varphi_\ell^*(\Phi_{\mathcal{X}}(E - dD)).$$

Expanding the left hand side, we find

$$\Phi_{\mathcal{X}^\ell}(E - dD) = \sum_{j=1}^{d} \Phi_{\mathcal{X}^\ell}(P_j - D).$$

On the other hand, as $E_{\mathcal{X}}$ avoids $T(\mathcal{X})$ we have that for each $j \in \{1, \ldots, d\}$ and for each non-archimedean place v the divisor $P_{j,\mathcal{X}}$ intersects precisely one component, so there is an index $i_j^v \in \{1, \ldots, r_v\}$ such that

$$\Phi_{\mathcal{X}}(E - dD) = \sum_{j=1}^{d} \sum_{v} \Phi_{\mathcal{X},v}\left(G^v_{i_j^v} - D\right).$$

See the beginning of Section 3 for the definition of $G^v_{i_j^v}$. Using Corollary 2.2, this implies

$$\sum_{j=1}^{d} \Phi_{\mathcal{X}^\ell}(P_j - D)^2 = \sum_{j=1}^{d} \sum_{v} \Phi_{\mathcal{X},v}\left(G^v_{i_j^v} - D\right)^2 = \sum_{j=1}^{d} \sum_{v} \left(\gamma_{D,v,i_j^v} - V_v(D)^2\right). \tag{10}$$

Because of (10) and by definition (6) of U_D we find

$$(E_{\mathcal{X}} \,.\, U_D) = \sum_{j=1}^{d} (P_{\mathcal{X},j} \,.\, U_D) = -\sum_{j=1}^{d} \gamma_{D,v,i_j^v} = -\sum_{j=1}^{d} \Phi_{\mathcal{X}^\ell}(P_j - D)^2 + dV_D^2. \tag{11}$$

This allows us to compare $h_{\overline{L_D}}$ to h_{NT}. We will use the Hodge Index Theorem due to Faltings and Hriljac (see for instance [9, §III.5]) which implies that if $P \in J$, then we have

$$h_{\mathrm{NT}}(P) \cdot [k : \mathbb{Q}] = -\left(\overline{\mathcal{O}}(Z_{\mathcal{X}}) \otimes \mathcal{O}(\Phi_{\mathcal{X}}(Z)) \,.\, \overline{\mathcal{O}}(Z_{\mathcal{X}})\right) = -\overline{\mathcal{O}}(Z_{\mathcal{X}})^2 + \mathcal{O}(\Phi_{\mathcal{X}}(Z))^2 \tag{12}$$

for any divisor Z of degree zero on X such that Z represents P.

We set $a' = -\overline{\mathcal{O}}(D_{\mathcal{X}})^2$. Then we get

$$\begin{aligned}
\sum_{i=1}^{d} h_{\mathrm{NT}}(j_D(P_i)) \cdot [k : \mathbb{Q}] &= \sum_{i=1}^{d} \left(-\overline{\mathcal{O}}\left(P_{j,\mathcal{X}^\ell} - D_{\mathcal{X}^\ell}\right)^2 + \mathcal{O}\left(\Phi_{\mathcal{X}^\ell}(P_j - D)\right)^2\right) \\
&= \sum_{j=1}^{d} \left(-\overline{\mathcal{O}}\left(P_{j,\mathcal{X}^\ell}\right)^2 + 2\left(\overline{\mathcal{O}}(P_{j,\mathcal{X}^\ell}) \,.\, \overline{\mathcal{O}}(D_{\mathcal{X}^\ell})\right) + \mathcal{O}\left(\Phi_{\mathcal{X}^\ell}(P_j - D)\right)^2\right) - d \cdot a' \\
&= \sum_{j=1}^{d} \left(\left(\overline{\mathcal{O}}(P_{j,\mathcal{X}^\ell}) \,.\, \overline{\omega}_{\mathcal{X}^\ell} \otimes \overline{\mathcal{O}}(2D_{\mathcal{X}^\ell})\right) + \mathcal{O}\left(\Phi_{\mathcal{X}^\ell}(P_j - D)\right)^2\right) - d \cdot a' \\
&= \left(\overline{\mathcal{O}}(E_{\mathcal{X}^\ell}) \,.\, \overline{\omega}_{\mathcal{X}^\ell} \otimes \overline{\mathcal{O}}(2D_{\mathcal{X}^\ell})\right) + \sum_{j=1}^{d} \mathcal{O}\left(\Phi_{\mathcal{X}^\ell}(P_j - D)\right)^2 - d \cdot a' \\
&= \left(\overline{\mathcal{O}}(E_{\mathcal{X}}) \,.\, \overline{\omega} \otimes \overline{\mathcal{O}}(2D_{\mathcal{X}})\right) - \left(\overline{\mathcal{O}}(E_{\mathcal{X}}) \,.\, \mathcal{O}(U_D)\right) - d \cdot a' + d \cdot \mathcal{O}(V_D)^2 \\
&= \left(\overline{\mathcal{O}}(E_{\mathcal{X}}) \,.\, \overline{L_D}\right) \\
&= d \cdot [k : \mathbb{Q}] \cdot h_{\overline{L_D}}(E).
\end{aligned}$$

Here the first equality holds by (12), the third equality holds because of the Arakelov adjunction formula (see [9, §IV.5]) and the fifth equality holds because of (9), (11) and because, by assumption, $E_{\mathcal{X}^l}$ does not intersect any vertical divisors contracted by φ_l.

The first assertion of the proposition is now immediate since the Néron-Tate height only takes nonnegative values, whence $h_{\overline{L_D}}(E) \geq 0$. The second assertion follows if we take $E = P$ in the argument above. □

Remark 4.7. Since the heights $h_{\overline{L_K}}(P)$ and $h_{\mathrm{NT}}([P-K])$ are related by Proposition 4.6, a comparison of the quantities $e'_{\overline{L_K}}$ and $a'(K)$ (cf. the interoduction of the present paper) which control the sharpness of our bound (1) and of Zhang's lower bound r, respectively, might be possible.

5. Semipositive hermitian line bundles – proof of Theorem 1.3

In this section we prove several quite general lemmas about certain hermitian line bundles on $\mathcal{X}$. All of these will be used in the proof of Theorem 1.3 in the case of $\overline{L_D}$. Several results of this section are quite similar to results from [16].

Definition 5.1. Let $\overline{L}$ be a hermitian line bundle on $\mathcal{X}$. We call a nonzero section s of $\overline{L}$ *effective* (resp. *strictly effective*) if $\|s\|_{\sup} \leq 1$ (resp. $\|s\|_{\sup} < 1$). We say that $\overline{L}$ is *ample* if L is ample and $H^0(\mathcal{X}, L^{\otimes n})$ has a basis consisting of strictly effective sections for $n \gg 0$. If this holds for $n = 1$, then we call $\overline{L}$ *very ample.*

Lemma 5.2. *Let $\overline{L}$ be a hermitian line bundle on $\mathcal{X}$. For any hermitian line bundle $\overline{M}$ on $\mathcal{X}$ and $a, b \in \mathbb{N}$ we set*

$$\overline{M}_{a,b} = \overline{M}^{\otimes a} \otimes \overline{L}^{\otimes b}.$$

If $\overline{L}^2 < 0$, then there exists no ample hermitian line bundle $\overline{M}$ on $\mathcal{X}$ with the following property: For all $a, b \in \mathbb{N}$ such that $\overline{M}_{a,b}^2 > 0$, we also have

$$\left(\overline{L} \,.\, \overline{M}_{a,b}\right) \geq 0. \tag{13}$$

Proof. This proof is somewhat similar to the first part of the proof of [16, Theorem 6.3]. Suppose that $\overline{L}^2 < 0$ and that $\overline{M}$ is a hermitian line bundle on $\mathcal{X}$ satisfying (13).

Since $\overline{M}$ is ample, it has positive arithmetic self-intersection by [16, Theorem 1.3]. Therefore, if $p(t)$ denotes the polynomial

$$p(t) = \overline{L}^2 + 2(\overline{L} \,.\, \overline{M})t + \overline{M}^2 t^2,$$

then there is a positive real number t_0 satisfying $p(t_0) = 0$ and $p(t) > 0$ for any $t > t_0$.

Now let $a, b \in \mathbb{N}$ such that $a/b > t_0$. Then we find

$$\overline{M}_{a,b}^2 = b^2 p(a/b) > 0.$$

By (13), we know that

$$\frac{1}{b}\left(\overline{L} \,.\, \overline{M}_{a,b}\right) = \overline{L}^2 + \frac{a}{b}(\overline{L} \,.\, \overline{M}) \geq 0.$$

In particular, our assumption that $\overline{L}^2 < 0$ implies

$$(\overline{L} \,.\, \overline{M}) > 0, \tag{14}$$

and also, since a/b can be arbitrary close to t_0,

$$\overline{L}^2 + (\overline{L} \,.\, \overline{M})t_0 \geq 0. \tag{15}$$

Now we can derive a contradiction as in the proof of [16, Theorem 6.3]. Namely, combining $p(t_0) = 0$ and (15) implies

$$(\overline{L} \,.\, \overline{M})t_0 + \overline{M}^2 t_0^2 \leq 0.$$

But this is impossible using $\overline{M}^2 > 0$ and (14). □

The following result provides us with a method to show that two hermitian line bundles on $\mathcal{X}$ have nonnegative intersection.

Lemma 5.3. *Let $\overline{L}$ and $\overline{M}$ be hermitian line bundles on $\mathcal{X}$. If $\overline{M}$ has an effective global section s such that $h_{\overline{L}}(\mathrm{div(s)}^{\mathrm{hor}}) \geq 0$ and $\overline{L}$ is relatively semipositive, then $(\overline{L} \,.\, \overline{M}) \geq 0$.*

Proof. According to [3, §3.2.2], we have

$$\left(\overline{L} \,.\, \overline{M}\right) = \frac{\left(\mathcal{O}(\mathrm{div(s)}^{\mathrm{ver}}) \,.\, \overline{L}\right)}{[k : \mathbb{Q}]} + h_{\overline{L}}(\mathrm{div(s)}^{\mathrm{hor}}) - \int \log \|s\|_{\sup} \, c_1(\overline{L}).$$

Now $\mathrm{div(s)}^{\mathrm{ver}}$ is effective, so the claim follows. □

Suppose we want to show that a hermitian line bundle $\overline{L}$ on $\mathcal{X}$ satisfies $\overline{L}^2 \geq 0$. Then using Lemma 5.2, it suffices to find some ample hermitian line bundle $\overline{M}$ on $\mathcal{X}$ such that under the assumption $\overline{L}^2 < 0$ we have $\left(\overline{L} \,.\, \overline{M}_{a,b}\right) \geq 0$ whenever $\overline{M}_{a,b}^2 > 0$.

If $\overline{L}$ is horizontally semipositive, then we can take any ample $\overline{M}$ and any effective section s of $\overline{M}_{a,b}^{\otimes n}$ (which exists for $n \gg 0$ by [16, Theorem 2.1], see [16, §8]) and apply Lemma 5.3. However, we want to apply Lemma 5.3 to $\overline{L} = \overline{L_D}$ which is not in general horizontally semipositive, but only $T(\mathcal{X})$-semipositive, so we have to be more careful with our choice of s.

Lemma 5.4 below tells us that, under the hypothesis $\overline{L}^2 < 0$, we can find $\overline{M}$ such that for $n \gg 0$ there are many effective sections of $\overline{M}_{a,b}$ whenever $\overline{M}_{a,b}^2 > 0$. Intuitively, it should be possible to find an effective section s avoiding a finite set of points if $\overline{M}_{a,b}$ has enough effective sections. Lemma 5.5 makes this intuition precise.

Lemma 5.4. *Suppose $\overline{L}$ is a relatively semipositive hermitian line bundle on $\mathcal{X}$ such that $\deg(L) > 0$ and $\overline{L}^2 < 0$. Then there exists an ample hermitian line bundle $\overline{M}$ on $\mathcal{X}$ such that for all $a, b \in \mathbb{N}$ satisfying*

$$\overline{M}_{a,b}^2 = \left(\overline{M}^{\otimes a} \otimes \overline{L}^{\otimes b}\right)^2 > 0,$$

the lattice $H^0\left(\mathcal{X}, M_{a,b}^{\otimes n}\right)$ has a basis consisting of strictly effective global sections for some $n = n(a, b) \gg 0$.

Proof. Let $\overline{M}$ be an ample hermitian line bundle on $\mathcal{X}$. We will scale $\overline{M}$ by $\alpha \in \mathbb{Q}_{\geq 0}$ such that the lemma holds for $\overline{M}(\alpha)$. Using [16, Theorem 1.5], it suffices to show that $\overline{M}_{a,b}(\alpha)^2 > 0$ implies that $\overline{M}_{a,b}(\alpha)$ is horizontally positive, since relative semipositivity is automatic.

Let $p_\alpha(t)$ denote the polynomial

$$p_\alpha(t) = \overline{L}^2 + 2\left(\overline{L} \,.\, \overline{M}(\alpha)\right)t + \overline{M}(\alpha)^2 t^2.$$

As in the proof of Lemma 5.2, there is some positive real number $t_0 = t_0(\alpha)$ such that $p_\alpha(t_0) = 0$ and $p_\alpha(t) > 0$ for all $t > t_0$.

Now let $m_L = -\inf_D h_{\overline{L}}(D)$ and $m_M = \inf_D h_{\overline{M}}(D)$, where we take the infima over all irreducible divisors D on X. If $m_L \geq 0$, then we can take $\alpha = 0$, so we suppose that $m_L < 0$.

Let D be some irreducible divisor on X. We will construct α such that

$$h_{\overline{M}_{a,b}(\alpha)}(D) \geq 0 \tag{16}$$

whenever $a/b > t_0(\alpha)$, which will prove the lemma.

Because of

$$\begin{aligned} h_{\overline{M}_{a,b}(\alpha)}(D) &= a h_{\overline{M}(\alpha)}(D) + b h_{\overline{L}}(D) \\ &\geq a m_M + a\alpha - b m_L, \end{aligned}$$

we need a nonnegative α such that $\frac{a}{b} \geq t_0(\alpha) \Rightarrow \frac{a}{b} \geq \frac{m_L}{\alpha + m_M}$, so α must satisfy

$$t_0(\alpha) \geq \frac{m_L}{\alpha + m_M}.$$

Hence (16) is easily seen to be equivalent to

$$(\alpha + m_M) r(\alpha) \geq m_L \left(\overline{M}^2 + 2\alpha \deg(M)\right) + (\alpha + m_M)\left((\overline{M} \,.\, \overline{L}) + \alpha \deg(L)\right), \tag{17}$$

where

$$r(\alpha) = \sqrt{\alpha^2 \deg(L)^2 + 2\alpha \deg(M)\left((\overline{M} \,.\, \overline{L}) - \overline{L}^2\right) + (\overline{M} \,.\, \overline{L})^2 - \overline{L}^2 \overline{M}^2}.$$

Note that $r(\alpha)$ is real since p_α always has real roots. Hence the left hand side of (17) is always nonnegative. If the right hand side is negative for some α, then (17) holds and we are done, so we may assume that the right hand side is also nonnegative.

We find that (17) holds if and only if $w(\alpha) \geq 0$, where w is a cubic polynomial in α, obtained by subtracting the square of the right hand side of (17) from the square of the left hand side. The leading coefficient of w is

$$-2 \deg(M)(\overline{L}^2 + 2 m_L \deg(L))$$

which is positive by our assumptions on $\overline{L}$. Hence (17) holds for $\alpha \gg 0$. □

Lemma 5.5. *Let $\overline{M}$ be an ample hermitian line bundle on $\mathcal{X}$ and let $P_1, \ldots, P_r$ be closed points on $\mathcal{X}$. Then for some $n \gg 0$ there exists a strictly effective global section s of $M^{\otimes n}$ such that $s(P_i) \neq 0$ for $i = 1, \ldots, r$.*

Proof. Without loss of generality we assume $\overline{M}$ is very ample. Since $\overline{M}$ is very ample as a hermitian line bundle, the lattice $H^0(X, M)$ is spanned by strictly effective sections $s_1, \ldots, s_m$. Moreover, since M is also very ample in the geometric setting, it is globally generated, i.e., there exists for each point P_i at least one section s_j such that $s_j(P_i) \neq 0$. Let $s_1, \ldots, s_k$ be a minimal set of such sections. Then for any even n the global section $s' = \sum_{j=1}^k s_j^n$ of M^n satisfies $s'(P_i) \neq 0$ for all $i = 1, \ldots, r$. But now for some even $n \gg 0$ the global section $s' = \sum_{j=1}^k s_j^n$ also satisfies the condition

$$\|s'\|_{\sup} \leq k \max_j \|s_j\|^n < 1.$$

□

Now let $D \in \mathrm{Div}_{\mathbb{Q}}(X)$ have degree 1. Recall Definition 3.1 of the hermitian line bundle $\overline{L_D}$ on $\mathcal{X}$. If $\overline{M}$ is a hermitian line bundle on $\mathcal{X}$ and a, b are positive integers, then we set

$$\overline{M}_{a,b} := \overline{M}^{\otimes a} \otimes \overline{L_D}^{\otimes b}.$$

We want to use Lemma 5.2 in order to prove Theorem 1.3, so we need to show that under the hypothesis $\overline{L_D}^2 < 0$ there is some hermitian line bundle $\overline{M}$ on $\mathcal{X}$ with positive self-intersection such that $\overline{M}_{a,b}^2 > 0$ implies $\left(\overline{L_D} \,.\, \overline{M}_{a,b}\right) \geq 0$.

Proposition 5.6. *Suppose that* $\overline{L_D}^2 < 0$ *and that* $D_{\mathcal{X}} \cap T(\mathcal{X}) = \emptyset$. *Then there exists an ample hermitian line bundle* $\overline{M}$ *on* $\mathcal{X}$ *such that for any positive integers* a, b *the following condition is satisfied: If* $\overline{M}_{a,b}^2 > 0$, *then there exists a positive integer* $n(a,b)$ *such that* $\overline{M}_{a,b}^{\otimes n(a,b)}$ *has an effective section* s *satisfying* $h_{\overline{L_D}}(\mathrm{div(s)}^{\mathrm{hor}}) \geq 0$.

Proof. It follows from Lemma 5.4 that there is an ample hermitian line bundle $\overline{M}$ on $\mathcal{X}$ such that $\overline{M}_{a,b}^2 > 0$ implies that $H^0\left(\mathcal{X}, M_{a,b}^{\otimes n}\right)$ has a basis consisting of strictly effective sections for n large enough. By [16, Theorem 1.3], $\overline{M}_{a,b}$ is ample.

We get from Lemma 5.5 that there is a multiple $n(a,b)$ of n and an effective section s of $\overline{M}_{a,b}^{\otimes n(a,b)}$ such that $\mathrm{div(s)}^{\mathrm{hor}}$ does not intersect the finite set $T(\mathcal{X}) \subset \mathcal{X}$. Using Proposition 4.6 we conclude $h_{\overline{L_D}}(\mathrm{div(s)}^{\mathrm{hor}}) \geq 0$. □

Now we can finally complete the proof of Theorem 1.3.

Proof of Theorem 1.3). Suppose that $\overline{L_D}^2 < 0$. Let $\overline{M}$ be as in Proposition 5.6 and let a, b be positive integers such that $\overline{M}_{a,b}^2 > 0$. It follows from Lemma 5.3 and Proposition 5.6 that we have

$$\left(\overline{L_D} \,.\, \overline{M}_{a,b}^{\otimes n(a,b)}\right) \geq 0$$

and thus

$$(\overline{L_D} \,.\, \overline{M}_{a,b}) \geq 0.$$

But by Lemma 5.2 this leads to a contradiction. □

6. Lower bounds on $\overline{\omega}^2$ – proof of Theorems 1.4 and 1.1

We finally get to our original problem, namely the derivation of lower bounds on $\overline{\omega}^2$. It follows from [5, Lemma A.1] that we can find a $\mathbb{Q}$-divisor $K \in \mathrm{Div}_{\mathbb{Q}}(X)$ such that $K_{\mathcal{X}} \cap T(\mathcal{X}) = \emptyset$ and such that $(2g-2) \cdot K$ is a canonical $\mathbb{Q}$-divisor on X. It is shown in [7] that

$$\mathcal{K} = (2g-2) \cdot (K_{\mathcal{X}} + V_K) \in \mathrm{Div}_{\mathbb{Q}}(\mathcal{X}) \tag{18}$$

is a canonical $\mathbb{Q}$-divisor on $\mathcal{X}$. By the latter we mean a $\mathbb{Q}$-divisor such that $\overline{\mathcal{O}}(\mathcal{K}) = \overline{\omega}$.

Proof of Theorem 1.4. Using (18) we get $K_{\mathcal{X}} = \frac{1}{2g-2} \cdot \mathcal{K} - V_K$. We can use this to rewrite $\overline{L_K}$ (cf. Definition 3.1) as

$$\overline{L_K} = \overline{\omega}^{\otimes \frac{g}{g-1}} \otimes \mathcal{O}(-2V_K - U_K) - 4g \cdot a,$$

where

$$a = \overline{\mathcal{O}}(K_{\mathcal{X}})^2 - \mathcal{O}(V_K)^2 = \frac{1}{4(g-1)^2}\overline{\omega}^2 - \frac{1}{g-1}(\overline{\omega} \,.\, \mathcal{O}(V_K)).$$

Hence we get

$$\overline{L_K}^2 = \frac{1}{g-1}\left(g\overline{\omega}^2 + (g-1)\mathcal{O}(2V_K + U_K)^2 - 2g(\overline{\omega} \,.\, \mathcal{O}(U_K))\right).$$

Since $\overline{L_K}^2 \geq 0$ by Theorem 1.3, the proof of Theorem 1.4 follows. □

Remark 6.1. We have developed the theory of $\overline{L_D}$ for rather general degree one divisors $D \in \mathrm{Div}_{\mathbb{Q}}(X)$. The main reason why we choose to work with K is that $K_{\mathcal{X}}$ has an obvious relation with $\overline{\omega}$. But there are other promising choices for D; for instance, we could take $D = \frac{1}{g^3-g}W$, where W is the divisor of Weierstrass points on X. This was suggested by Ariyan Javanpeykar. In fact, it is easy to see that the divisor $\mathcal{V}$ used in [8, Lemma 5.1] to extend W to a divisor on $\mathcal{X}$ with good properties is a valid choice for our V_W, so it would be interesting to study properties of $\overline{L_D}$.

Next we use Theorem 1.4 to prove Theorem 1.1.

Proof of Theorem 1.1. From Proposition 3.4 we know that the assumptions of Theorem 1.4 are satisfied. Therefore it suffices to show that $(\overline{\omega} \,.\, \mathcal{O}(U_K)) \geq 0$. To this end, we will use the adjunction formula.

Let v be a non-archimedean place of k and let $\Gamma_1, \ldots, \Gamma_r$ denote the irreducible components of $\mathcal{X}_v$. The adjunction formula and minimality imply that we have

$$a_i = (\mathcal{K} \,.\, \Gamma_i) = -\Gamma_i^2 + 2p_a(\Gamma_i) - 2 \geq 0.$$

for all i. As before, we let $G_i \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ be a $\mathbb{Q}$-divisor on $X \times k_v^{\mathrm{nr}}$ of degree one such that $(G_{i,\mathcal{X}} \,.\, \Gamma_j) = \delta_{ij}$.

It is easy to see that if $E_1, \ldots, E_n \in \mathrm{Div}_{\mathbb{Q}}(X \times k_v^{\mathrm{nr}})$ have degree one and $E = \frac{1}{n}(E_1 + \ldots + E_n)$, then

$$U_E = \frac{1}{n}\left(U_{E_1} + \ldots + U_{E_n}\right).$$

Hence we may assume that $K \times k_v^{\mathrm{nr}} = G_t$ for some $t \in \{1, \ldots, r\}$.

By definition of $U_K = \sum_v U_{K,v}$, we have

$$(U_{K,v} \,.\, \mathcal{K}) = \sum_{i=1}^{r} -V_v(G_i)^2 a_i + 2\sum_{i=1}^{r}(V_v(G_i) \,.\, V_v(G_t))a_i.$$

Therefore it certainly suffices to show

$$\sum_{i=1}^{r}(V_v(G_i) \,.\, V_v(G_t))a_i = 0. \tag{19}$$

Recall the definition of $V_v(G_i) = \sum_{j=1}^{r} c_j(G_i)\Gamma_j$. Let $a'_j := \frac{a_j}{2g-2}$ and

$$w_j(G_i) = a'_j - (G_{i,\mathcal{X}} \,.\, \Gamma_j) = a'_j - \delta_{ij}.$$

If $w(G_i) = \left(w_j(G_i)\right)_j \in \mathbb{Q}^r$ and $c(G_i) = \left(c_j(G_i)\right)_j \in \mathbb{Q}^r$, then we have

$$c(G_i) = M^+ \cdot w(G_i), \tag{20}$$

where M^+ is, as usual, the Moore-Penrose pseudoinverse of minus the intersection matrix of $\mathcal{X}_v$. Now we compute

$$\begin{aligned}(V_{G_i} \,.\, V_{G_t}) = \left(V_{G_i} \cdot \sum_{j=1}^{r} c_j(G_t)\Gamma_j \right) &= \sum_{j=1}^{r} c_j(G_t) \cdot (V_{G_i} \,.\, \Gamma_j) \\ &= \sum_{j=1}^{r} c_j(G_t) w_j(G_i)\end{aligned}$$

Using (20) we find

$$\sum_{i=1}^{r} (V_v(G_i) \,.\, V_v(G_t)) a_i = -(2g-2) w(G_t)^T \cdot M^+ \cdot \sum_{i=1}^{r} w(G_i) \cdot a_i'$$

and thus the proof of (19) follows from the fact that for each $j \in \{1, \ldots, r\}$ we get

$$\sum_{i=1}^{r} w_j(G_i) a_i' = a_j' \sum_{i=1}^{r} a_i' - a_j' = 0,$$

since $\sum_{i=1}^{r} a_i' = 1$. □

Remark 6.2. If all multiplicities are simple, then we can rewrite the lower bound (1) above using the entries of the matrices $M_{\mathcal{X}_v}$ and $M^+_{\mathcal{X}_v}$ and the numbers $a_{v,i}$, all defined in Section 2. This allows for rapid computations of the lower bound (1) for specific arithmetic surfaces.

Let v be a non-archimedean place of k. For simplicity, we will only present a formula for the intersection on the right hand side of the inequality in Theorem 1.4 above v. To get a formula for the lower bound (1) one only needs to sum over all v. We also omit all references to v on the right hand side.

A long and tedious computation shows that for any $\mathbb{Q}$-divisor D of degree 1 whose closure on $\mathcal{X}$ avoids $T(\mathcal{X})$ the following equality

$$\begin{aligned}&(\mathcal{O}(U_D) + 2V_D)_v^2 + 2(\mathcal{K} \,.\, \Gamma_i)_v \\ &= \; 4\sum_{i=1}^{r} v_i(D) n_{ii} - 4\sum_{i=1}^{r}\sum_{j=1}^{r}\sum_{k=1}^{r} v_i(D) n_{ij} n_{kk} m_{jk} - \sum_{i=1}^{r}\sum_{j=1}^{r} n_{ii} n_{jj} m_{ij} \\ &\quad + 4g \sum_{i=1}^{r}\sum_{j=1}^{r} a_i' a_j' n_{ij} - 4g \sum_{i=1}^{r} a_i' n_{ii}\end{aligned}$$

holds. Note that this is not independent of D; however, one can show that as long as there exists a component Γ_i such that $(D_{\mathcal{X}} \,.\, \Gamma_j) = \delta_{ij}$, the bound is independent of i. In particular, if we know that there exists a canonical divisor of the form $(2g-2)P$, where $P \in X(k)$, then we can compute our lower bound directly using only the matrices M_v, M_v^+ and the vectors $(a'_{j,v})_j$ for all v of bad reduction.

For such X an explicit comparison between our bound and Zhang's lower bound r is possible without constructing V_K or U_K, leading to a strategy for verifying the effective Bogomolov conjecture for X (see Section 1).

7. Fermat curves of prime exponent

In Section 6 we derived a nontrivial lower bound on $\overline{\omega}^2$ for minimal arithmetic surfaces with simple multiplicities. In this section we compute lower bounds on $\overline{\omega}^2$ in a situation where Theorem 1.1 is not applicable, namely for minimal regular models of Fermat curves of prime exponent. Along the way, we construct U_K and show that $\overline{L_K}$ is relatively semipositive.

We start with a brief review of the notation from [7]. Let $p > 3$ be a prime number, let

$$F_p : X^p + Y^p = Z^p$$

denote the Fermat curve with exponent p over $k = \mathbb{Q}(\zeta_p)$, where ζ_p is a primitive p-th root of unity. Let $\mathcal{F}_p^{\min}$ denote the minimal regular model of F_p over $\mathbb{Z}[\zeta_p]$ as computed by McCallum (cf. [12]). We denote the components of the only non-reduced special fiber of $\mathcal{F}_p^{\min}$ by $L_i, i \in I$, where

$$I = \{x, y, z,\ \alpha_1,\ \alpha_{1,1}, \ldots, \alpha_{1,p},\ \alpha_2,\ \alpha_{2,1}, \ldots, \alpha_{2,p},\ \ldots,\ \alpha_r,\ \alpha_{r,1}, \ldots, \alpha_{r,p},\ \beta_1, \ldots, \beta_s\}$$

and $2r + s = p - 3$ (see [7]). Since a picture of the configuration of the non-reduced special fiber of $\mathcal{F}_p^{\min}$ is not too helpful, we give in Figure 1 the configuration of the special fiber of a non-minimal model $\mathcal{F}_p$ such that contracting the unique exceptional component L on $\mathcal{F}_p$ yields $\mathcal{F}_p^{\min}$. See [12, 7] for details.



Figure 1. The configuration of the special fiber of $\mathcal{F}_p$ above v. All components have genus 0. The only component with self-intersection number -1 is L.

Note that L_{α_i} has multiplicity two. Therefore Proposition 3.4 and Theorem 1.1 do not apply and we have to show relative semipositivity of $\overline{L_K}$ and nontriviality of the bound from Theorem 1.4 directly for a suitable $\mathbb{Q}$-divisor K.

Since there is only one place v of bad reduction, we will omit it from the notation for the sake of simplicity.

It was shown by Curilla and Kühn in [7] that a canonical divisor of F_p is given by $(2g-2)\cdot K$, where $K = S_x$ for a certain k-rational point S_x on F_p whose Zariski closure $\mathcal{S}_x$ intersects only L_x. In order to compute a lower bound on $\overline{\omega}^2$ using the results of the previous sections we will find U_K. This means that we first have to compute V_{G_i}, where for each $i \in I$ the Zariski closure $G_{i,\mathcal{X}\times\mathcal{O}_v^{\mathrm{nr}}}$ of the divisor $G_i \in \mathrm{Div}(X \times k_v^{\mathrm{nr}})$ of degree one satisfies $(G_{i,\mathcal{X}\times\mathcal{O}_v^{\mathrm{nr}}} \,.\, L_j) = \delta_{ij}$.

Lemma 7.1. *We have*

- $V_{G_i} = \frac{1}{p} L_i$, *if* $i \in \{x, y, z, \beta_j\}$,
- $V_{G_{\alpha_i}} = \frac{1}{p} L_{\alpha_i} + \frac{1}{2p} \sum_{j=1}^{p} L_{\alpha_{ij}}$,
- $V_{G_{\alpha_{i,j}}} = -\frac{1}{p} L_{\alpha_i} + \left(\frac{1}{2} - \frac{1}{2p}\right) L_{\alpha_{ij}} - \frac{1}{2p} \sum_{k \neq j} L_{\alpha_{ik}}$.

Proof. Recall that the divisor V_{G_i} must satisfy

$$(V_{G_i} \,.\, L_j) = a'_j - (G_{i, \mathcal{X} \times \mathcal{O}_v^{\mathrm{nr}}} \,.\, L_j) \tag{21}$$

for all $j \in I$, where $a'_j = \frac{1}{2g-2}(-L_j^2 + 2p_a(L_j) - 2)$. Since all components have genus zero and self-intersection $1-p$ except for the components $L_{\alpha_{ij}}$, which have genus zero and self-intersection -2, we get $a'_j = 0$ for the latter and $a'_j = \frac{1}{p}$ for all other components. Checking the validity of the Lemma reduces to checking (21) for each $i \in I$ which is a simple computation that we leave to the reader.

Note that for V_{G_x} this was essentially shown in [7, Proposition 8.3]. □

Corollary 7.2. *A canonical divisor on* $\mathcal{F}_p^{\min}$ *is given by* $(2g-2)\mathcal{S}_x + \frac{1}{p} L_x$ *and we have*

$$\begin{aligned} U_K =& \frac{1-p}{p^2} L_x + \frac{1+p}{p^2} \left(L_y + L_z + \sum_{j=1}^{s} L_{\beta_j} \right) + \frac{1+p/2}{p^2} \sum_{i=1}^{r} L_{\alpha_i} \\ &+ \frac{p^2/2 + p/2 - 3}{p^2} \sum_{i=1}^{r} \sum_{j=1}^{p} L_{\alpha_{ij}} \end{aligned}$$

Proof. The first assertion follows immediately from (18), the fact that $(2g-2)S_x$ is a canonical divisor on F_p and Lemma 7.1.

For the second assertion, recall the definition of U_K from Section 3:

$$U_K = \sum_{i \in I} \left(2(V_{G_i} \,.\, V_{S_x}) - V_{G_i}^2 \right) L_i.$$

Next we compute

$$\begin{aligned} V_{G_i}^2 &= \left(\frac{1}{p} L_i \right)^2 = \frac{1}{p^2}(1-p) \text{ for } i \in \{x, y, z, \beta_j\} \\ (V_{G_i} \,.\, V_{S_x}) &= \left(\frac{1}{p} L_i \,.\, \frac{1}{p} L_x \right) = \frac{1}{p^2} \text{ for } i \in \{y, z, \beta_j\} \\ (V_{G_{\alpha_i}} \,.\, V_{S_x}) &= \left(\frac{1}{p} L_{\alpha_i} + \frac{1}{2p} L_{\alpha_{ij}} \,.\, \frac{1}{p} L_x \right) = \frac{1}{p^2} \end{aligned}$$

$$\begin{aligned}
(V_{G_{\alpha_{ij}}} \,.\, V_{S_x}) &= \left(-\frac{1}{p}L_{\alpha_i} + \left(\frac{1}{2} - \frac{1}{2p}\right)L_{\alpha_{ij}} - \frac{1}{2p}\sum_{k\neq j} L_{\alpha_{ik}} \,.\, \frac{1}{p}L_x\right) = -\frac{1}{p^2} \\
(V_{G_{\alpha_i}})^2 &= \left(\frac{1}{p}L_{\alpha_i} + \frac{1}{2p}L_{\alpha_{ij}}\right)^2 = \frac{1}{p^2}(1-p) + \frac{p}{p^2} - \frac{2p}{4p^2} = \frac{1}{p^2} - \frac{1}{2p} \\
(V_{G_{\alpha_{ij}}})^2 &= \left(-\frac{1}{p}L_{\alpha_i} + \left(\frac{1}{2} - \frac{1}{2p}\right)L_{\alpha_{ij}} - \frac{1}{2p}\sum_{k\neq j} L_{\alpha_{ik}}\right)^2 \\
&= \frac{1}{p^2}(1-p) - \frac{1}{2}\left(1-\frac{1}{p}\right)^2 - \frac{1}{2p^2}(p-1) - \frac{1}{p}\left(1-\frac{1}{p}\right) + \frac{p-1}{p^2} \\
&= \frac{1}{2p^2}(2-p-p^2)
\end{aligned}$$

A simple computation proves the corollary. □

Lemma 7.3. *The hermitian line bundle $\overline{L_K}$ is relatively semipositive.*

Proof. Recalling Definition 3.1 of $\overline{L_K}$ we see that we need to show

$$a_i + 2(\mathcal{S}_x \,.\, L_i) - (U_K \,.\, L_i) \geq 0 \tag{22}$$

for all $i \in I$, where $a_i = 0$ for $i = \alpha_{ij}$ and $a_i = p-3$ for all other components.

As usual, we will distinguish between the different components L_i to prove (22). Throughout, we will use that $s = p - 3 - 2r$ to eliminate s. We start with L_x and find

$$(U_K \,.\, L_x) = \frac{1-p}{p^2}(1-p) + (s+2)\frac{1+p}{p^2} + r\frac{1+p/2}{p^2} = 2 - \frac{2}{p} - \frac{r}{p^2} - \frac{3r}{2p}$$

which gives

$$a_x + 2(\mathcal{S}_x \,.\, L_x) - (U_K \,.\, L_x) \geq p - 3 > 0.$$

Let $i \in \{y, z, \beta_j\}$. Then we get

$$(U_K \,.\, L_i) = \frac{1-p}{p^2} + \frac{1+p}{p^2}(1-p) + (s+1)\frac{1+p}{p^2} + r\frac{1+p/2}{p^2} = -\frac{2}{p} - \frac{r}{p^2} - \frac{3r}{2p}$$

and thus

$$a_i + 2(\mathcal{S}_x \,.\, L_i) - (U_K \,.\, L_i) \geq p - 3 > 0.$$

It remains to consider the components L_{α_i} and $L_{\alpha_{ij}}$. The computations are similar to the ones above, but tedious and hence are omitted. The upshot is that we get inequalities

$$a_{\alpha_i} + 2(\mathcal{S}_x \,.\, L_{\alpha_i}) - (U_K \,.\, L_{\alpha_i}) = p - 3 - \frac{1}{2}p + 1 + \frac{5}{p} + \frac{r}{p^2} + \frac{3r}{2p} \geq \frac{p}{2} - 2 > 0$$

and

$$a_{\alpha_{ij}} + 2(\mathcal{S}_x \,.\, L_{\alpha_{ij}}) - (U_K \,.\, L_{\alpha_{ij}}) = 1 + \frac{1}{2p} - \frac{7}{p^2} > 0.$$

This shows that (22) is satisfied for all components L_i which proves the lemma. □

Proposition 7.4. *We have*

$$\begin{aligned}\overline{\omega}^2 \geq & \frac{1}{4p^3(p-1)(p-2)}\big((4+2r)p^6-(32+10r)p^5+(10+19r)p^4+(124\\ & -r-25r^2)p^3+(-56+52r+31r^2)p^2+(156-328r+112r^2)p\\ & +144-24r+60r^2\big)\log p\end{aligned}$$

Proof. By Theorem 1.4 and Lemma 7.3 we know that

$$\overline{\omega}^2 \geq -\frac{g-1}{g}\mathcal{O}(2V_K+U_K)^2+2(\overline{\omega}\,.\,\mathcal{O}(U_K)).$$

Hence we need to compute the terms on the right hand side.

First note that

$$\begin{aligned}2V_K+U_K=\frac{1+p}{p^2}&\left(L_x+L_y+L_z+\sum_{j=1}^{s}L_{\beta_j}\right)\\ &+\frac{1+p/2}{p^2}\sum_{i=1}^{r}L_{\alpha_i}+\frac{p^2/2+p/2-3}{p^2}\sum_{i=1}^{r}\sum_{j=1}^{p}L_{\alpha_{ij}}\end{aligned}$$

Using this, it is not hard to verify that

$$\begin{aligned}(2V_K+U_K)^2 =\; & -\frac{pr}{2}-\frac{r}{2}+1+\frac{1}{p}(\frac{7}{4}r-5)+\frac{1}{p^2}(\frac{25}{4}r^2-5r-1)\\ & +\frac{1}{p^3}(17-30r+11r^2)+\frac{1}{p^4}(12-2r+5r^2).\end{aligned} \tag{23}$$

For the computation of $(\mathcal{K}\,.\,U_K)$, where $\mathcal{K}$ is a canonical divisor on $\mathcal{F}^{\min}$, we use the adjunction formula. Namely, if δ_i denotes the multiplicity of L_i in U_K, then we have

$$(\mathcal{K}\,.\,U_K)=\sum_{i\in I}\delta_i a_i$$

and hence

$$(\mathcal{K}\,.\,U_K)=(p-3)\left(\frac{1-p}{p^2}+(s+2)\frac{1+p}{p^2}+r\frac{1+p/2}{p^2}\right). \tag{24}$$

Combining (23) and (24) proves the proposition after a little algebra. □

The proof of Theorem 1.5 is now immediate.

Proof. (of Theorem 1.5) For the general lower bound, use Proposition 7.4 and $0 \leq r \leq \frac{1}{2}p-\frac{3}{2}$. To derive the lower bounds for $p=5,7$, we use that $r=0$ if $p=5$ and $r=2$ if $p=7$. □

Remark 7.5. The upper bound computed by Curilla and Kühn in [7] is of order $\mathcal{O}(gp\log p)$, i.e. of order $\mathcal{O}(p^3\log p)$. Hence it seems that with our methods we cannot derive an asymptotic formula for $\overline{\omega}^2$.

References

[1] A. Abbes and E. Ullmo, *Auto-intersection du dualisant relatif des courbes modulaires* $X_0(N)$, J. Reine Angew. Math. **484**, 1–70 (1997).

[2] S. Bosch, W. Lütkebohmert and M. Raynaud, *Néron models*, Springer-Verlag, Berlin (1990).

[3] J.-B. Bost, H. Gillet and C. Soulé, *Heights of projective varieties and positive Green forms*, J. Am. Math. Soc. **7**, No. 4, 903–1027 (1994).

[4] J.-F. Burnol, *Weierstrass points on arithmetic surfaces*, Invent. Math. **107**, 421–432 (1992).

[5] T. Chinburg and R. Rumely, *The capacity pairing*, J. Reine Angew. Math. **434**, 1–44 (1993).

[6] Z. Cinkir, *The tau constant and the discrete Laplacian matrix of a metrized graph*, Eur. J. Comb. **32**, No. 4, 639–655 (2011).

[7] C. Curilla and U. Kühn, *On the arithmetic self-intersection number of the dualizing sheaf for Fermat curves of prime exponent*, Preprint (2009). arXiv:math/0906.3891v1 [math.NT]

[8] R. de Jong, *Arakelov invariants of Riemann surfaces*, Doc. Math. **10**, 311–329 (2005).

[9] S. Lang, *Introduction to Arakelov theory*, Springer-Verlag, New York (1988).

[10] Q. Liu, *Stable reduction of finite covers of curves*, Compositio Math. **142**, 101–118 (2006).

[11] H. Mayer, *Self-Intersection of the Dualizing Sheaf of Modular Curves* $X_1(N)$, PhD Thesis, HU Berlin (2011).

[12] W.G. McCallum, *The degenerate fiber of the Fermat curve*, in *Number theory related to Fermat's last theorem*, Progr. Math. **26**, Birkhäuser, Boston, 57–70 (1982).

[13] A. Moriwaki, *Lower bounds of self-intersection of dualizing sheaves on arithmetic surfaces with reducible fibres*, Math. Ann. **305**, 183–190 (1996).

[14] X. Sun, *On relative canonical sheaves of arithmetic surfaces*, Math. Z. **223**, 709–723 (1996).

[15] E. Ullmo, *Positivité et Discrétion des Points Algébriques des Courbes*, Ann. Math. (2) **147** (1), 167–179 (1998).

[16] S. Zhang, *Positive line bundles on arithmetic surfaces*, Ann. Math. (2) **136**, 569–587 (1992).

[17] S. Zhang, *Admissible pairing on a curve*, Invent. Math. **112**, 171–193 (1993).